



TAROLA PLASTIC SURGERY  
NICHOLAS A. TAROLA, MD

### **Breast Lift**

**Remove Wrap** the day following surgery.

**You may shower after you remove the wrap.** If you have steri strips on your incision, leave them in place and pat dry after showering.

**Wear a soft cup or sports bra for 4 weeks after your surgery.** You may re-wrap yourself in ace wrap if it is more comfortable until seen at your first post-operative visit.

**No underwire bra until cleared by Dr. Tarola or his staff (usually after 4 weeks).**

**No heavy lifting for 4-6 weeks after surgery.** You may resume normal activities in 2 weeks.

**Tightness in the chest region and stiffness:** Tingling, burning, or intermittent shooting pain is normal as the skin, tissues, and sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. If you have drains, you may experience additional localized discomfort.

**Consistent sharp pain should be reported to our office immediately.**

**Hypersensitivity of nipples or lack of sensitivity:** This is normal and will gradually resolve over time. You may also experience a small amount of fluid or milk seeping through the nipples. If this becomes painful or excessive, notify our office immediately.

**Shiny skin or any itchy feeling:** Swelling can cause the breast skin to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling of the breasts. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. If the skin becomes red and hot to the touch, contact our office immediately.

**Asymmetry, the breasts look different, or heal differently:** Breasts may look or feel quite different from one another in the days following surgery. This is normal. No two breasts in nature or following surgery are perfectly symmetrical.

**CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:**

- **A high fever (over 101°), severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations**
- **Any pain that cannot be controlled by your pain medication**
- **Bright red skin that is hot to the touch**
- **Excessive bleeding or fluid seeping through the incisions**
- **A severely misshapen breast or bruising that is localized to one breast or region of the chest**

**To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the treated region.** Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses, for no longer than 20-minute intervals.